

Short Communication

Status and distribution of the Near Threatened Tibetan argali *Ovis ammon hodgsoni* in Ladakh, India: effect of a hunting ban

TSEWANG NAMGAIL, JOSEPH L. FOX and YASH VEER BHATNAGAR

Abstract The Near Threatened Tibetan argali *Ovis ammon hodgsoni* is distributed across the Tibetan Plateau and its peripheral mountains. Within India it occurs in Ladakh (Jammu and Kashmir) and Sikkim, and the population was estimated to be c. 200 in the early 1990s. Hunting of the species was banned in Jammu and Kashmir in the 1980s but the effect of this hunting moratorium on the population has not previously been assessed. We conducted surveys in the proposed Gya-Miru Wildlife Sanctuary and the neighbouring Tsokar Basin, areas reported to have a relatively high abundance of argali within Ladakh. We also opportunistically surveyed other areas and collected secondary information about the species' occurrence in other parts of Ladakh. A total of 127 animals were counted during the surveys. Based on this number and other small populations reported earlier by us and others, we estimate a total population of 300–360 argali in Ladakh. Although past population estimates were approximate, the present estimate, which includes areas not previously surveyed, suggests there has been no substantial change in the population of argali in Ladakh since the early 1980s. Factors other than hunting therefore appear to be impeding argali population recovery in this region. Pashmina-producing goats are the most abundant livestock within the argali's range and, owing to the recent increase in demand for this fibre, the goat population is increasing and this may be hindering the recovery of the argali.

Keywords Gya-Miru, India, Ladakh, *Ovis ammon hodgsoni*, Tibetan argali, Tsokar.

The Tibetan argali *Ovis ammon hodgsoni* is a wild sheep of the subfamily Caprinae (Bovidae). Although widely distributed over large mountainous tracts of Central Asia,

O. a. hodgsoni is in a critical condition with <7,000 individuals left in the wild (Schaller, 1998). In India the estimated population in the 1990s was c. 200, with most occurring in Ladakh (Fox et al., 1991a) and a few in Sikkim (Fox & Johnsingh, 1997). It is listed as a fully protected (Schedule I) species under the Indian Wildlife (Protection) Act of 1972. The Tibetan argali is one of only two subspecies of *O. ammon* listed on Appendix I of CITES (UNEP-WCMC, 2008), and *O. ammon* is categorized as Near Threatened on the IUCN Red List (IUCN, 2008).

The range of the Tibetan argali includes the Tibetan Plateau and its marginal mountains, encompassing c. 2.5 million km² (Shackleton, 1997), but its population is highly fragmented throughout this vast range (Fox et al., 1991a; Schaller, 1998). Argali were the least encountered wild ungulate during extensive surveys on the Tibetan Plateau in the late 1980s and early 1990s (Schaller, 1998). The population in Ladakh is mostly confined to rolling mountains of eastern areas bordering China (Fox et al., 1991a) but its precise range is unknown. Using habitat suitability modelling Chundawat & Qureshi (1999) identified c. 10,988 km² in Ladakh as potential habitat for argali. Although argali are present in isolated parts of this area (M. Ranjitsinh, pers. comm.) Bhatnagar & Wangchuk (2001) did not observe the species in large tracts of potential range but did locate small populations in three previously unreported areas (Skagzung, Phobrang and Chumur).

Trophy hunting is thought to have contributed to the earlier decimation of the argali population in Ladakh (Ward, 1924; Fox et al., 1991a), as the animal has large horns coveted by hunters (Adair, 1899). The last trophy hunting for the subspecies in Ladakh occurred in 1975 in the Tsabra catchment of the proposed Gya-Miru Wildlife Sanctuary (Namgail et al., 2004a). Nomadic herders, army personnel and some Government officers also hunted the species for meat, and the argali's preference for open areas, often close to human settlements, made it especially vulnerable (Namgail et al., 2004b). Hunting of argali was banned in India in the early 1980s but whether this has led to a population recovery has not previously been evaluated.

We carried out surveys during March–April 2003 in the Gya-Miru Wildlife Sanctuary and the neighbouring Tsokar Basin (Main survey area, Fig. 1), areas known to support a relatively high abundance of argali in Ladakh (Fox et al., 1991a). Additionally, we searched for argali in other potential

TSEWANG NAMGAIL* (Corresponding author) and YASH VEER BHATNAGAR International Snow Leopard Trust (India Program), Nature Conservation Foundation, 3076/5, IV Cross, Gokulam Park, Mysore 570 002, Karnataka, India. E-mail namgail@ncf-india.org

JOSEPH L. FOX Department of Biology, University of Tromsø, Norway.

*Also at: Resource Ecology Group, Department of Environmental Sciences, Wageningen University, Droevendaalsesteeg 3a, 6708 PB Wageningen, The Netherlands.

Received 25 February 2008. Revision requested 7 April 2008.

Accepted 20 May 2008.




Fig. 1 Current distribution of the Tibetan argali in the Ladakh Trans-Himalaya of India. The Line of Actual Control (LAC) is the effective border between India and China and the Line of Control (LoC) is the effective border between India and Pakistan. The rectangle on the inset indicates the location of the main figure.

areas (Bhatnagar & Wangchuk, 2001; Namgail, 2001). We also compiled published data and gathered secondary information from knowledgeable wildlife officials, paramilitary personnel, wildlife researchers and local pastoralists about the

occurrence and numbers of argali in areas not covered by surveys.

Using topographic features such as ridges and streams the main survey area was partitioned into six localities:

Puyul, Gya, Kyam-Tsabenama, Tsabra, Tisaling-Shiabuk, and catchments in and around the the Gya-Miru Wildlife Sanctuary and Tsokar Basin. On average we spent 17 hours (over 2 days) surveying each area (on foot, although horses were sometimes used to reach a site). From trails and ridge lines, slopes were scanned with binoculars and spotting scopes. Once a group of argali was located we recorded the time, date, group size and sex/age composition. Males were classified, on the basis of horn size, as class I (2–3 years old), II (3–5), III (5–7) and IV (≥ 7 ; Fedosenko et al., 1995). Counts were repeated on 2 consecutive days in all six areas. The possibility of double counting on the same day as well as on consecutive days was prevented by not including any group that was similar to a previously observed group in its size, sex and age composition. Size of surveyed areas were determined from a 1:250,000 digitized map of the survey area using the geographical information system *ArcMap v. 9.2* (ESRI, Redmond, California). Argali density was calculated by dividing the total number counted in each block (the higher count of the 2 days) by area. Areas that were not visible from the survey trails or were unusable by argali, such as lakes and glaciers, were excluded.

A total of 127 argali were located in three of the six areas surveyed (Table 1). Mean group size was $7.6 \pm \text{SE } 0.97$ (range = 1–29). Of the 127 counted, 48 were seen in the Tsabra catchment, with 15 males, 20 females, four yearlings and nine lambs. The sex ratio was thus 75:100 in favour of females, and the lamb:ewe ratio was 45:100. There was a preponderance of class III (40%, $n = 48$) and IV (26%) males.

We counted 20 argali in the Hemis National Park in the summer of 2000 (Namgail, 2001) and this population still survives (Rinchen Wangchuk, pers. comm.). In July 2004 we observed a group of three argali in the upper Hanle Valley from where, according to local informants, they were thought to have gone extinct c. 20 years earlier. Interviews with local people and paramilitary forces in the adjacent Kuyul Valley suggested there is a population of 10–15 argali near Demchok, apparently moving between

India and China. Although we did not observe any argali in the Chumur Valley during a visit in 2004, interviews with paramilitary personnel suggested that there is a population of 10–20, close to the border with China.

During a 1-week survey in January 2007 around Phobrang and Marsemik La, a mountain pass leading into the Chhang Chhenmo Valley, we saw two males, three females, one yearling and one lamb. Conversations with wildlife officials, local nomadic herders and researchers, however, suggested that 100–130 argali occur in and around the Chhang Chhenmo Valley, historically an important site for argali. Although a small population of 5–10 individuals has previously been reported from the Yaya Tso area (Raghuandan S. Chundawat, pers. comm.), we did not observe any argali during our surveys there in July 2008. According to local informants 10–20 argali survive in Kharnak and c. 15 in the Skazung area (Fig. 1).

Based on our surveys and secondary information we estimate that the total argali population in Ladakh is 300–360. This includes small subpopulations in areas not previously surveyed and therefore suggests there has been no substantial change in the population of argali in Ladakh since the late 1980s. Given that trophy hunting halted > 20 years ago (Rauf Zargar, Wildlife Warden, pers. comm.) we conclude that, although sporadic poaching for meat might have occurred, factors other than hunting are hindering the recovery of the argali population in Ladakh.

Ladakh is currently undergoing substantial socio-economic change, and populations of wild ungulates such as argali are being affected by development activities that influence habitat use (Bhatnagar et al., 2006; Namgail et al., 2007b). Pashmina (cashmere wool) is a mainstay of the economy of people living within the argali's range, and livestock populations, especially those of cashmere-producing goats, have increased substantially (Namgail et al., 2007a, 2008). Studies on the interaction between livestock and argali have shown that livestock grazing not only depletes resources required by argali but also physically displaces argali from productive pastures (Namgail et al., 2007b). Increasing livestock populations, and disturbances created by livestock grazing and collateral activities, are thus the likely factors hindering the recovery of argali. Livestock grazing has negative influences on populations of other argali subspecies in Central Asia (Harris & Bedunah, 2001; Maroney, 2005; Wingard, 2005; Schaller & Kang, 2008).

We conclude that the ban on argali hunting has not been sufficient to protect and enhance argali populations in Ladakh. The relatively high abundance of argali in the Gya-Miru Wildlife Sanctuary and the adjacent Tsokar Basin suggests this area may be the best location in which to protect argali in India but even there the livestock population is increasing because of demand for pashmina. Initial conservation efforts should focus on reducing pressure on the argali population, possibly by freeing some

TABLE 1 Terrain type and area of the six localities surveyed in the proposed Gya-Miru Wildlife Sanctuary and neighbouring Tsokar Basin (Main survey area, Fig. 1), with estimates of the number of argali counted (see text for further details) and argali density.

Locality	Terrain type	Area (km ²)	No. counted	Density (km ⁻²)
Puyul	Rugged	37	0	
Gya	Moderate	30	0	
Kyam-Tsabenama	Moderate	47	0	
Tsabra	Moderate	60	48	0.8
Tisaling-Shiabuk	Undulating	134	15	0.2
Tsokar Basin	Undulating	164	64	0.4
<i>Total</i>		472	127	0.3*

*Calculated only for the blocks where argali were observed

areas from livestock grazing, as has been done for the Tibetan gazelle *Procapra picticaudata* in Ladakh (Namgail et al., 2008). The argali population in the Gya-Miru Wildlife Sanctuary and Tsokar Basin could serve as a source population to recolonize adjacent areas, as apparently happened in the late 1970s in Hemis National Park (Fox et al., 1991b).

We are initiating a dialogue with the people of Gya-Miru about the possibility of creating an Argali Reserve devoid of livestock grazing. Although there is also a small population of argali in Sikkim (Fox & Johnsingh, 1997) its concentration at the international border, apparently moving back and forth between Indian and Chinese-controlled territory (Chanchani et al., in press), makes it difficult to assure unilaterally its conservation within India. Thus the Gya-Miru and Tsokar areas of Ladakh appear currently to be the areas best suited for the conservation of argali in India.

Acknowledgements

The study was funded by the University of Tromsø-Wildlife Institute of India Institutional Co-operation Program, the Rufford Small Grants Foundation and the Wildlife Conservation Society. Jigmet Takpa and Salim Ul-Haq helped in acquiring the research permit. Lobzang and Tsewang Morup helped in the field and R. Raghunath prepared the map.

References

- ADAIR, F.E.S. (1899) *A Summer in High Asia*. Willian Clowes and Sons, London, UK.
- BHATNAGAR, Y.V., MISHRA, C. & WANGCHUK, R. (2006) Decline of the Tibetan gazelle in Ladakh. *Oryx*, 40, 229–232.
- BHATNAGAR, Y.V. & WANGCHUK, R. (2001) Status survey of large mammals in eastern Ladakh and Nubra. In *Conservation Biodiversity in the Trans-Himalaya: New Initiatives for Field Conservation in Ladakh*, pp. 108–125. Wildlife Institute of India, Dehradun, India.
- CHANCHANI, P., RAWAT, G.S. & GOYAL, S.P. (in press) Status and distribution of trans-Himalayan ungulates in Tso Lhamo, Sikkim, India. *Oryx*.
- CHUNDAWAT, R.S. & QURESHI, Q. (1999) *Planning Wildlife Conservation in Leh and Kargil Districts of Ladakh, Jammu & Kashmir*. Wildlife Institute of India, Dehradun, India.
- FEDOSENKO, A.K., WEINBERG, P.I. & VALDEZ, R. (1995) Argali sheep survey in the headwaters of the Korunduk River, Kirghizstan. *Mammalia*, 59, 452–455.
- FOX, J.L. & JOHNSINGH, A.J.T. (1997) Country report for India. In *Wild Sheep and Goats and their Relatives: Status Survey and Conservation Action Plan for Caprinae* (ed. D.M. Shackleton), pp. 215–231. IUCN, Gland, Switzerland and Cambridge, UK.
- FOX, J.L., NURBU, C. & CHUNDAWAT, R.S. (1991a) The mountain ungulates of Ladakh, India. *Biological Conservation*, 58, 167–190.
- FOX, J.L., NURBU, C. & CHUNDAWAT, R.S. (1991b) Tibetan argali (*Ovis ammon hodgsoni*) establish a new population. *Mammalia*, 55, 448–451.
- HARRIS, R.B. & BEDUNAH, D.J. (2001) *Sheep vs Sheep: Argali and Livestock in Western China*. Report to National Geographic Society, Washington, DC, USA.
- IUCN (2008) *2008 IUCN Red List of Threatened Species*. IUCN, Gland, Switzerland. [Http://www.iucnredlist.org](http://www.iucnredlist.org) [accessed 20 January 2009].
- MARONEY, R.L. (2005) Conservation of argali *Ovis ammon* in western Mongolia and the Altai-Sayan. *Biological Conservation*, 121, 231–241.
- NAMGAIL, T. (2001) *Habitat selection and ecological separation between sympatric Tibetan argali and blue sheep in northern India*. MPhil thesis, University of Tromsø, Tromsø, Norway.
- NAMGAIL, T., BAGCHI, S., MISHRA, C. & BHATNAGAR, Y.V. (2008) Distributional correlates of the Tibetan gazelle *Procapra picticaudata* in Ladakh, northern India: towards a recovery programme. *Oryx*, 42, 107–112.
- NAMGAIL, T., BHATNAGAR, Y.V., MISHRA, C. & BAGCHI, S. (2007a) Pastoral nomads of the Indian Changthang: production system, land use and socio-economic changes. *Human Ecology*, 35, 497–504.
- NAMGAIL, T., FOX, J.L. & BHATNAGAR, Y.V. (2004a) *Argali and Livestock: Friends or Foes? Interactions between Tibetan Argali and Domestic Sheep/Goats in a Prospective Trans-Himalayan Reserve*. Report to the International Snow Leopard Trust, Seattle, USA.
- NAMGAIL, T., FOX, J.L. & BHATNAGAR, Y.V. (2004b) Habitat segregation between sympatric Tibetan argali *Ovis ammon hodgsoni* and blue sheep *Pseudois nayaur* in the Indian Trans-Himalaya. *Journal of Zoology*, 262, 57–63.
- NAMGAIL, T., FOX, J.L. & BHATNAGAR, Y.V. (2007b) Habitat shift and time budget of the Tibetan argali: the influence of livestock grazing. *Ecological Research*, 22, 25–31.
- SCHALLER, G.B. (1998) *Wildlife of the Tibetan Steppe*. Chicago University Press, Chicago, USA.
- SCHALLER, G.B. & KANG, A. (2008) Status of Marco Polo sheep *Ovis ammon polii* in China and adjacent countries: conservation of a Vulnerable subspecies. *Oryx*, 42, 100–106.
- SHACKLETON, D.M. (ed.) (1997) *Wild Sheep and Goats and their Relatives: Status Survey and Conservation Action Plan for Caprinae*. IUCN, Gland, Switzerland, and Cambridge, UK.
- UNEP-WCMC (2008) *CITES Species Database*. [Http://www.cites.org/eng/resources/species.html](http://www.cites.org/eng/resources/species.html) [accessed 20 January 2009].
- WARD, A. (1924) The mammals and birds of Kashmir and the adjacent hill provinces. *Journal of the Bombay Natural History Society*, 29, 879–887.
- WINGARD, J.R. (2005) *Dietary Overlap Between Argali Sheep and Domestic Livestock in Mongolia*. MSc thesis, University of Montana, Missoula, USA.

Biographical sketches

TSEWANG NAMGAIL and YASH VEER BHATNAGAR are studying mountain ungulates, pastoralism and human-wildlife conflicts in the Indian Trans-Himalaya. They work closely with local communities, finding mutually acceptable ways to mitigate conflicts between wildlife and pastoralists, and innovative ways for the long-term conservation of threatened species such as the snow leopard and its prey. JOSEPH L. FOX has been studying wildlife and pastoralism on the Tibetan Plateau and its peripheral areas, including Ladakh, for over 3 decades. He has a particular interest in snow leopard conservation in central Asia and also wide experience in reconciling wildlife conservation and livestock production in Tibet and various Himalayan countries.