

Potential distribution of and priority conservation areas for the Endangered Caatinga howler monkey *Alouatta ululata* in north-eastern Brazil

ROBÉRIO FREIRE FILHO and JORGE M. PALMEIRIM

Abstract The Caatinga of north-eastern Brazil is the largest and most diverse seasonally dry tropical forest in the Americas and is home to numerous endemic species. However, only 1.2% of the area is under full protection, and given the ongoing decline of this biome there is an urgent need to expand the protected area network. The Endangered Caatinga howler monkey *Alouatta ululata* is almost endemic to the Caatinga, and is a potential umbrella species for the protection of its biodiversity. Using all available distribution data and our own surveys we applied *Maxent* and *Zonation* spatial modelling to identify the range of *A. ululata*, and priority conservation areas for the species, maximizing habitat quality and connectivity, and minimizing conservation constraints. The top 10% priority areas cover 34,400 km² and mostly coincide with good remnants of Caatinga. Only priority areas in the northern part of the species' range are protected, so it is essential to create new protected areas in the centre and south of the range. *Maxent* modelling indicates that the species depends on good tree cover, but even inside protected areas we observed recent deforestation, illustrating the urgency to improve management. *Maxent* also indicated that aridity limits the species' range, and therefore the ongoing aridification of the Caatinga is a threat to its future. The protection of *A. ululata* requires establishing new protected areas in priority locations and improving management of existing protected areas. Preservation of priority areas for the Caatinga howler monkey also represents an opportunity for the conservation of other important biodiversity in the region.

Keywords *Alouatta ululata*, Brazil, Caatinga, Caatinga howler monkey, *Maxent*, primate, species distribution modelling, *Zonation*

Supplementary material for this article is available at <https://doi.org/10.1017/S0030605318001084>

ROBÉRIO FREIRE FILHO (Corresponding author) Programa de Pós-Graduação em Biologia Animal, Universidade Federal de Pernambuco, Centro de Biociências, Departamento de Zoologia, Av. Prof. Moraes Rego, 1235, Cidade Universitária, Cep. 50670-420, Recife-Pernambuco, Brazil. E-mail freirefilho@outlook.com

JORGE M. PALMEIRIM Departamento de Biologia Animal, Centro de Ecologia, Evolução e Alterações Ambientais, Faculdade de Ciências, Universidade de Lisboa, Lisboa, Portugal

Received 15 February 2018. Revision requested 17 April 2018.
Accepted 7 August 2018.

Introduction

The Caatinga covers c. 735,000 km² of north-eastern Brazil (Leal et al., 2005) and is considered to be a significant wilderness area (Aguiar et al., 2002). It is the largest and most diverse seasonally dry tropical forest in the Americas and harbours large numbers of endemic species (DRYFLOR et al., 2016). However, its natural vegetation has been declining at an alarming rate (Beuchle et al., 2015) as a result of land-use intensification (Aguiar et al., 2002; Leal et al., 2005). There is an urgent need for conservation measures to protect the Caatinga, with only 1.2% of the area currently under full protection (DRYFLOR et al., 2016). Protected areas need to be expanded, and charismatic species, such as large primates, can facilitate this process (Ducarme et al., 2013) as they function as umbrella species for the conservation of valuable but more discreet biodiversity within their range. One of the species with the greatest potential for this role is the Caatinga howler monkey *Alouatta ululata*, which requires vast areas of suitable habitat to maintain viable populations.

The Caatinga howler monkey is categorized as Endangered on the IUCN Red List because of its small and declining population, a consequence of tree cover loss, habitat fragmentation and hunting (Oliveira & Kierulff, 2008). Most of the species' range is within the Caatinga, although it extends into the Cerrado (Oliveira & Kierulff, 2008), but its limits are poorly known, which is a major constraint for the planning of conservation measures (Oliveira & Kierulff, 2008).

Species distribution modelling is a tool for mapping geographical distributions and studying how they are influenced by environmental variables (Miller, 2010). It is widely applied in conservation science and its models may support the selection of areas for conservation (Araújo et al., 2002). The most common approach in species distribution modelling is maximum-entropy modelling, often applied using *Maxent* software (Phillips et al., 2006).

The identification of priority areas for conservation of species is a fundamental step in developing conservation plans (Pressey et al., 2007), and computational tools have been developed to carry out this process of prioritization, taking into consideration factors such as habitat quality, connectivity and conservation cost (Kukkala & Moilanen, 2013). *C-Plan* (Pressey et al., 2009), *Marxan* (Watts et al., 2009) and *Zonation* (Moilanen et al., 2005) are examples of approaches and packages developed for conservation prioritization. The general objective of all these packages is




FIG. 1 Locations of records and reports of the Caatinga howler monkey *Alouatta ululata*, and interviews with members of the local rural community, throughout the species' known range, in north-eastern Brazil.

similar, although their strategies vary. For example, *Zonation*, used in this study, prioritizes landscapes by iteratively removing the least valuable remaining areas while accounting for connectivity and generalized complementarity (Moilanen et al., 2011).

The overarching objective of this project was to conduct spatially explicit analyses that contribute to the planning of measures to conserve *A. ululata* and the ecosystems with which it is associated. Our specific objectives were to (1) carry out field surveys to collect information on the distribution of the species, (2) identify areas where further surveys are needed, (3) develop a model to generate a potential species distribution map, and examine the environmental determinants of the species' distribution, (4) identify the areas with most potential for the conservation of the species, and (5) determine the degree of coverage of the priority areas by existing protected areas. We use our results to make spatially explicit recommendations for actions needed to improve the conservation of *A. ululata* and of the many species that depend on the same Caatinga habitats.

Study area

The study area includes the known range of *A. ululata*, across the states of Maranhão, Piauí and Ceará (Fig. 1). In the spatial analysis we included not only the area encompassing all the known locations of the species, but also a 30-km wide buffer zone around this. The aim was to identify areas that may be suitable for the species but that are outside its currently known range.

Methods

Data collection

We compiled existing information on the distribution of *A. ululata*, most of which was collected during 2004–2010 by the National Center for Research and Conservation of Brazilian Primates (CPB/ICMBio). As the number of direct observations by researchers is low (20 records), we also used reports obtained in CPB/ICMBio interviews with members of the local rural community during 2004–2010 (112 reports). Interview-based distribution analysis can complement direct monitoring data in the case of easily identifiable species (Anadón et al., 2010; Brittain et al., 2018). We also carried out surveys in two regions for which information was scarce, during August 2016–May 2017 (Fig. 1). We interviewed 112 farmers and hunters who lived or worked close to areas with natural vegetation. To minimize bias we did not reveal that *A. ululata* was the focus of our interviews (Freire Filho et al., 2018). We asked about other mammals present in the region before asking about *A. ululata* (Freire Filho et al., 2018). All interviews were conducted with the consent of the participants.

Some records based on interviews were initially referenced with the coordinates of the place of the interview, usually farmhouses. We replaced these coordinates with those of the nearest area of natural environment within a 2-km radius of the original coordinates (approximately the distance up to which the vocalization of the species can be heard). Points without natural environment within a 2-km radius were excluded. This procedure adds locational uncertainty, but *Maxent* modelling can make useful predictions even

TABLE 1 Predictive variables used in our analysis of the potential distribution of and priority conservation areas for the Caatinga howler monkey *Alouatta ululata*.

Variable	Description	Source
Maxent variables		
% tree cover	Canopy closure for all vegetation > 5 m height	Global forest change 2000–2014; Hansen et al. (2013)
Aridity index	Rainfall deficit for potential vegetative growth (higher values represent greater aridity)	CGIAR Consortium for Spatial Information; Zomer et al. (2008)
Bio17	Precipitation of driest quarter	WorldClim 1.4; Hijmans et al. (2005)
Bio15	Precipitation seasonality (SD of monthly precipitation expressed as %)	WorldClim 1.4; Hijmans et al. (2005)
Forest canopy height	Global 1 km forest canopy height	SDAT (2011), Simard et al. (2011)
Ruggedness index	Measurement of terrain heterogeneity generated in QGIS using SRTM3 data	USGS (2000), Riley et al. (1999)
Constraint layers		
Anthropogenic areas	Land cover map from GlobCover project; all categories (croplands & urban areas) were joined in a single class	ESA (2009)
Influence of roads	Buffer of road influence 18 km either side; this distance was selected subjectively by visually analysing land cover along roads in the study area	IBGE (2015)
Population density	Global Rural–Urban Mapping Project, Version 1 (GRUMPv1): Population Density Grid	SEDAC (2011), Balk et al. (2006)

when the occurrence data include a moderate level of locational error (Graham et al., 2008). Our initial database included 184 occurrences, 52 from our surveys and 132 from CPB/ICMBio. However, to minimize problems associated with spatial biases in sampling we used spatial filtering (Kramer-Schadt et al., 2013), reducing the number of occurrences in oversampled areas by using only one within a radius of 5 km. This filtering procedure reduced the number of occurrences used in the modelling to 117 (20 direct observations and 70 reports from CPB/ICMBio, and one observation and 26 reports from our surveys).

Modelling of potential distribution

We used Maxent to identify variables influencing the distribution of *A. ululata* and generate a distribution map (Phillips et al., 2006). The choice of environmental variables was guided by the species' biology; *A. ululata* is arboreal, feeds on leaves, fruits and other plant parts, and lives in a semi-arid region influenced by seasonal rainfall and high temperatures. We expected areas with higher precipitation to be more suitable during the most critical period of the year, the dry season. Furthermore, we hypothesized that tree cover and tree height would influence suitability, and that areas with rugged terrain would be more suitable because terrain ruggedness tends to be an obstacle to habitat destruction and to provide some protection from hunting. With the help of a matrix of correlations between all candidate variables, we selected a set of six predictors (Table 1)

that were not highly correlated ($|r| < 0.70$; Rainho & Palmeirim, 2013) and that were biologically meaningful.

Prior to running Maxent all layers were converted to the WGS 1984 geographical coordinate system and to a cell size of 30 arc seconds (c. 1 km²), using IDRISI Selva (Eastman, 2012) and QGIS 2.8 (QGIS Development Team, 2016). In Maxent we used the following settings: convergence threshold (10–5), maximum iterations (500), regularization multiplier (1), maximum number of background points (104), linear, quadratic, product and hinge features, random seed generation and 50 replicates. The resulting map is in logistic format, with the probability of presence for each cell being in the range 0–100% (Phillips, 2008). To select a suitability threshold for the potential distribution map we used the methodology described in Rainho & Palmeirim (2013), which facilitates the selection of the smallest area including most occurrences. The area selected was that corresponding to the Maxent suitability threshold 70%, which encompassed 83% of the occurrences (Fig. 2); above this threshold the inclusion of more occurrences would force the addition of a disproportionately large area.

Prioritizing areas for conservation

We used Zonation (Moilanen et al., 2005) to prioritize areas for the conservation of *A. ululata*. Zonation generates a priority map that can be used to inform decision-making. Cost efficiency can be considered in this prioritization through the inclusion of a cost layer. We assume that conservation




FIG. 2 Relationship between per cent probability of presence and per cent of presence points for the Caatinga howler monkey, used to define the suitability threshold for the species in the *Maxent* model.




FIG. 3 Probability of presence of the Caatinga howler monkey, as predicted by the *Maxent* model.

cost efficiency is higher in areas with fewer anthropogenic constraints, and thus generated a single constraints layer combining three thematic layers: human population density, proximity to roads, and anthropogenic areas (i.e. urban areas and farmland). These three layers (Table 1) were given equal weight in the generation of the constraints layer. *Zonation* assigned this layer negative weights and combined it with the map of potential distribution of *A. ululata* generated with *Maxent*.

In *Zonation* we selected distribution smoothing as the aggregation method. It considers fragmentation to be undesirable and thus retains areas that are well interconnected. The size of the smoothing kernel used was 6 km, a value

based on the distances crossed by various *Alouatta* species outside their usual home ranges (Glander, 1992; Crockett, 1998). As a cell removal rule we used core-area zonation, which is appropriate when importance is given to core areas, i.e. locations with the highest suitability in terms of abundance or high probability of occurrence (Moilanen et al., 2017). To avoid losing valuable areas and to keep structural connectivity we selected the options ‘add edge points’ and ‘edge removal’ (Moilanen et al., 2017).

To identify the priority regions lacking protection, we overlaid the *Zonation* map with existing protected areas. We used a map of forest loss during 2000–2014 (Hansen et al., 2013) to identify recent deforestation in priority areas.

TABLE 2 Results of the *Maxent* distribution model for *A. ululata*.

Variables	% contribution	Permutation importance	Training gain without	Training gain with only	Test gain without	Test gain with only	AUC* without	AUC* with only
% tree cover	55.411	51.928	0.789	0.502	0.714	0.475	0.814	0.751
Bio17	15.786	16.139	1.009	0.145	0.909	0.149	0.846	0.661
Aridity index	10.516	17.884	1.043	0.363	0.949	0.359	0.847	0.727
Ruggedness index	7.102	1.778	1.025	0.140	0.966	0.120	0.854	0.622
Bio15	5.179	7.624	1.025	0.041	0.907	0.042	0.845	0.578
Forest canopy height	6.004	4.644	1.031	0.204	0.985	0.169	0.849	0.663

*AUC, area under the curve.




FIG. 4 The relationship of the probability of presence of the Caatinga howler monkey with six environmental variables: (a) per cent tree cover, (b) aridity, (c) precipitation of driest quarter, (d) Ruggedness, (e) precipitation seasonality, and (f) forest canopy height.

Results

The mean test area under the curve (AUC) for the *Maxent* distribution model of *A. ululata* was $0.857 \pm \text{SD } 0.032$, indicating a high efficiency in distinguishing presence from random background locations (Fig. 3). The variables with greatest contributions in the potential distribution model were per cent tree cover, precipitation of driest quarter (Bio17) and aridity index (Table 2). The jackknife analysis corroborates these results (Table 2). Probability of presence increases with tree cover, canopy height and terrain ruggedness (Fig. 4). In the case of the climatic variables (aridity, precipitation in driest quarter and precipitation seasonality) the highest probability of occurrence tends to be in the intermediate values (Fig. 4).

Results of the *Zonation* prioritization are shown in Fig. 5a. For graphical clarity we show only two levels of priority: high priority (the best 20%) and top priority (the best 10%, c. 34,400 km²). These priority areas can be divided into four ecologically distinct regions: (1) mangroves, in the mouth of the Parnaíba River; (2) enclaves, encompassing the humid areas of north-west Ceará and northern Piauí; (3) Caatinga, a vast region within the Caatinga biome; and (4) border, located along the border of the states of Piauí and Maranhão (Fig. 5a).

Circa 21% of the most important areas for the conservation of *A. ululata* are within legally protected areas (Fig. 5a). However, the coverage of priority regions by these protected areas is uneven; important parts of mangrove and enclave




FIG. 5 (a) Priority areas for the conservation of the Caatinga howler monkey *A. ululata* in mangrove, enclave, Caatinga and border areas, and existing protected areas; (b) forest loss during 2000–2014 within priority areas; (c) areas requiring survey work, new protected areas, and improved management.

areas have some level of legal protection, whereas border and Caatinga areas are virtually unprotected (Table 3).

Although the loss of tree cover during 2000–2014 was scattered, it occurred throughout the high-priority areas, including in protected areas. In some enclaves of northern Piauí deforestation is more widespread and there are larger deforested patches (Fig. 5b). Overall c. 3.8% of prime area for the conservation of *A. ululata* was deforested during 2000–2014 (Table 3).

Discussion

Potential distribution

A visual analysis shows a good correspondence between the distribution and density of the occurrence of the species with the map generated by *Maxent*. However, there are suitable areas without occurrences, possibly because of local extinctions (e.g. as a result of hunting) or a lack of survey effort (Oliveira & Kierulff, 2008). There are also a few occurrences

TABLE 3 Protected area and forest loss in each of the four priority regions identified for *A. ululata*.

	Inclusion in protected areas		Forest loss (2000–2014)	
	%	Area (km ²)	%	Area (km ²)
Mangrove	69	2,359.028	8	272.640
Enclave	50	9,814.051	4	776.058
Caatinga	10	63.619	2	926.061
Border	3	15,519.234	7	1,039.788
<i>All four regions</i>	21	27,755.932	4	3,014.507

in areas with low *Maxent* suitability, which can be explained by the extensive loss of natural habitat (Oliveira & Kierulff, 2008).

An isolated population inhabits a small humid enclave in Acopiara, Ceará, separated from the range of the species by > 100 km of unsuitable dry Caatinga (Fig. 1; Oliveira et al., 2007). It is probably a remnant of a broader distribution of the species when humid forest dominated the region

(Carmignotto et al., 2012), but could have been introduced by people long ago (monkeys are often kept as pets by Indigenous People, so this population could potentially have originated from escaped animals).

The eastern limit of the range of *A. ululata* is clear, defined by the high-aridity conditions in central Ceará. However, the western limit is ill defined and possibly explained by the wet conditions prevailing in Maranhão. Moreover, competition with *Alouatta belzebul*, a closely related species occurring further west, may contribute to shaping the western limit of the range of *A. ululata*. It is unclear if the two taxa are distinct species or differentiated populations of the same species (Viana et al., 2015).

The variables used in the *Maxent* model are coherent with the biology of the species (Oliveira & Kierulff, 2008). Per cent tree cover was the most important of these variables and suitability was low up to c. 50% tree cover, which is probably explained by the marked arboreal habit of the species (Oliveira & Kierulff, 2008). The species' diet of leaves and fruit may also contribute to its dependence on dense tree cover, as trees are its main food source (Oliveira & Kierulff, 2008). Moreover, the results indicate that *A. ululata* prefers tall forest, even though it occupies a variety of woodland types (Oliveira & Kierulff, 2008).

The second and third most influential variables in the model are both climatic and related: aridity index and precipitation in the driest quarter (Bio17). The results indicate that regions with an extreme dry season are unsuitable for *A. ululata*, suggesting vulnerability to the ongoing aridification of the Caatinga (Torres et al., 2017), where it is predicted that by the end of the 21st century temperatures may be up to 3.5–4.5 °C higher and rainfall 40–50% lower than at present (PBM, 2013).

Priority areas for conservation and current level of conservation

The priority areas identified using *Zonation* (Fig. 5a) should combine high habitat suitability, identified by *Maxent*, with good connectivity and low conservation constraints. This should make our results a good basis for conservation planning. However, both *Maxent* and *Zonation* modelling are affected by sources of error, such as inaccuracies in species occurrences and environmental layers, which create uncertainty (Graham et al., 2008; Moilanen et al., 2017). New and better models should be generated as more information on species or better environmental layers become available, and management decisions may have to be adjusted to take the new results into account. However, the inevitable uncertainty associated with models should not be an obstacle to their careful use in planning conservation action; in the face of rapid environmental change the risks of inaction are probably greater than those of using models judiciously (Wiens et al., 2009).

The priority areas identified encompass ecologically distinct regions. For example, the *A. ululata* population that inhabits the mangroves at the mouth of the Parnaíba River has a unique ecology (e.g. a diet composed of mangrove plants). The groups that inhabit the enclave region of Ceará live mostly in humid areas with open ombrophilous forest. Adaptations to these different environments may have resulted in populations of *A. ululata* with distinct behaviours, ecologies, gene pools and even morphologies. It is desirable to protect the various ecological contexts in which the species is present. Virtually all priority areas with protected status are in the northern part of the species' range. The south, including the region with most occurrences, is unprotected. This situation reflects the scarcity of protected areas in the Caatinga (de Marques & Peres, 2015; DRYFLOR et al., 2016).

The level of conservation provided by most protected areas in the range of *A. ululata* is minimal. Of the nine relevant protected areas only two (Ubajara and Sete Cidades National Parks) are fully protected, with nature preservation being their main objective, and only indirect use of natural resources allowed (MMA, 2016). All other conservation units allow sustainable use of natural resources (MMA, 2016), which needs to be well managed to avoid damage. However, management is insufficient because of a lack of human and financial resources. Moreover, in this region awareness of the protected areas is low (Drummond et al., 2009).

Our analysis indicated that deforestation is ongoing throughout most of the priority areas, even within protected areas (Fig. 5b), in line with the general trend of tree cover loss in the Caatinga (Beuchle et al., 2015). The situation is better only in federally protected National Parks, where we did not identify recent deforestation. Thus it is evident that protected areas currently make only a small contribution to the protection of *A. ululata* and the other natural values of the Caatinga.

Implications for conservation

The *Maxent* analysis indicates that good tree cover and levels of aridity lower than those prevailing in the Caatinga region are critical for *A. ululata*, suggesting that the species is affected by both deforestation and climate change. Unfortunately, tree cover is declining (Beuchle et al., 2015), even in protected areas (Fig. 5c), and arid conditions are increasing (Torres et al., 2017). This negative context should be taken into consideration in planning the conservation of the species.

Based on the results of our spatial analyses we recommend four conservation actions for *A. ululata*. These actions vary across the range of the species (Fig. 5c).

Firstly, *Maxent* modelling indicates that the range of *A. ululata* includes poorly surveyed areas where the presence of the species has not been confirmed, and the *Zonation* analysis indicates that some of these areas have a high level of

conservation priority. These areas should be surveyed as a matter of urgency (Fig. 5c).

Secondly, our results demonstrate that the legal protection of areas suitable for *A. ululata* is uneven. The central and southern parts of the species' range, which include some of the potentially best areas for its conservation, are not currently protected. Thus the designation of new state, federal and private protected areas is critical, especially in the larger contiguous priority areas (Fig. 5c).

Thirdly, and concurrently, management of the existing protected areas requires improvement.

Fourthly, knowledge on the biology of *A. ululata* is still insufficient to plan effective management measures and therefore, in addition to survey work to clarify the status of the species, further research is required on aspects of the species' ecology that are critical to its conservation.

The priority areas identified by the *Zonation* models mostly coincide with good-quality remnants of Caatinga, which are also important for the rich biodiversity of this biome, threatened by habitat destruction but poorly covered by protected areas (DRYFLOR et al., 2016). Given the species' medium body size, the conservation of viable populations of *A. ululata* requires the maintenance of large areas of well-preserved habitat, making it a good umbrella species with potential to contribute to the protection of the biodiversity in this unique ecoregion.

Acknowledgements This study was funded by a Rufford Foundation grant (19646-1). RFF is supported by a scholarship from Fundação de Amparo à Ciência e Tecnologia de Pernambuco (IBPG-1236-2.05/16). We thank the rural communities who collaborated in this research; Thieres Pinto, Gabriela Linhares, Andressa Fraga and Nádia Freitas for assistance in the field; and Marcos Fialho and Leandro Jerusalinsky (CPB-ICMBio) for providing presence records of *A. ululata*.

Author contributions Study conception and design, writing: RFF, JMP; field work and data analysis: RFF.

Conflicts of interest None.

Ethical standards This research complies with the *Oryx* Code of Conduct. No specimens were killed or collected. Interviews were conducted in accordance with the ethical standards of the British Sociological Association (BSA, 2017).

References

- AGUIAR, J., LACHER, T.E. & DA SILVA, J.M.C. (2002) The Caatinga. In *Wilderness: Earth's Last Wild Places* (eds R.A. Mittermeier, C.G. Mittermeier, P. Robles Gil, J. Pilgrim, G.A.B. da Fonseca, T. Brooks & W.R. Konstant), pp. 174–181. Cemex, Mexico City, Mexico.
- ANADÓN, J.D., GIMÉNEZ, A. & BALLESTAR, R. (2010) Linking local ecological knowledge and habitat modelling to predict absolute species abundance on large scales. *Biodiversity and Conservation*, 19, 1443–1454.
- ARAÚJO, M.B., WILLIAMS, P.H. & FULLER, R.J. (2002) Dynamics of extinction and the selection of nature reserves. *Proceedings of the Royal Society of London B: Biological Sciences*, 269, 1971–1980.
- BALK, D.L., DEICHMANN, U., YETMAN, G., POZZI, F., HAY, S.I. & NELSON, A. (2006) Determining global population distribution: methods, applications and data. *Advances in Parasitology*, 62, 119–156.
- BEUCHLE, R., GRECCHI, R.C., SHIMABUKURO, Y.E., SELIGER, R., EVA, H.D., SANO, E. & ACHARD, F. (2015) Land cover changes in the Brazilian Cerrado and Caatinga biomes from 1990 to 2010 based on a systematic remote sensing sampling approach. *Applied Geography*, 58, 116–127.
- BRITAIN, S., NGO BATA, M., DE ORNELLAS, P., MILNER-GULLAND, E. & ROWCLIFFE, M. (2018) Combining local knowledge and occupancy analysis for a rapid assessment of the forest elephant *Loxodonta cyclotis* in Cameroon's timber production forests. *Oryx*, published online 19 March 2018.
- BSA (BRITISH SOCIOLOGICAL ASSOCIATION) (2017) *BSA Statement of Ethical Practice*. https://www.britisoc.co.uk/media/24310/bsa_statement_of_ethical_practice.pdf [accessed 12 September 2018].
- CARMIGNOTTO, A.P., VIVO, M.D. & LANGGUTH, A. (2012) Mammals of the Cerrado and Caatinga: distribution patterns of the tropical open biomes of central South America. In *Bones, Clones, and Biomes: The History and Geography of Recent Neotropical Mammals* (eds B.D. Patterson & L.P. Costa), pp. 307–350. University of Chicago Press, Chicago, USA.
- CROCKETT, C.M. (1998) Conservation biology of the genus *Alouatta*. *International Journal of Primatology*, 19, 549–578.
- DE MARQUES, A.A.B. & PERES, C.A. (2015) Pervasive legal threats to protected areas in Brazil. *Oryx*, 49, 25–29.
- DUCARME, F., LUQUE, G.M. & COURCHAMP, F. (2013) What are “charismatic species” for conservation biologists? *BioSciences Master Reviews*, 10, 1–8.
- DRUMMOND, J.A., FRANCO, J.L.A. & NINIS, A.B. (2009) Brazilian federal conservation units: a historical overview of their creation and of their current status. *Environment and History*, 15, 463–491.
- DRYFLOR, BANDA-R, K., DELGADO-SALINAS, A., DEXTER, K.G., LINARES-PALOMINO, R., OLIVEIRA-FILHO, A. et al. (2016) Plant diversity patterns in neotropical dry forests and their conservation implications. *Science*, 353, 1383–1387.
- EASTMAN, J.R. (2012) *IDRISI Selva Manual Version 17.01*. Clark University, Worcester, USA.
- ESA (EUROPEAN SPACE AGENCY) (2009) *GlobCover 2009 (Global Land Cover Map)*. http://due.esrin.esa.int/page_globcover.php [accessed 12 September 2018].
- FREIRE FILHO, R., PINTO, T. & BEZERRA, B.M. (2018). Using local ecological knowledge to access the distribution of the Endangered Caatinga howler monkey (*Alouatta ululata*). *Ethnobiology and Conservation*, 7, 1–22.
- GLANDER, K.E. (1992) Dispersal patterns in Costa Rican mantled howling monkeys. *International Journal of Primatology*, 13, 415–436.
- GRAHAM, C.H., ELITH, J., HIJMANS, R.J., GUIBAN, A., TOWNSEND PETERSON, A., LOISELLE, B.A. & THE NCEAS PREDICTING SPECIES DISTRIBUTIONS WORKING GROUP (2008) The influence of spatial errors in species occurrence data used in distribution models. *Journal of Applied Ecology*, 45, 239–247.
- HANSEN, M.C., POTAPOV, P.V., MOORE, R., HANCHER, M., TURBANANOVA, S.A., TYUKAVINA, A. et al. (2013) High-resolution global maps of 21st-century forest cover change. *Science*, 342, 850–853.
- HIJMANS, R.J., CAMERON, S.E., PARRA, J.L., JONES, P.G. & JARVIS, A. (2005) Very high resolution interpolated climate surfaces for global land areas. *International Journal of Climatology*, 25, 1965–1978.

- IBGE (INSTITUTO BRASILEIRO DE GEOGRAFIA E ESTATÍSTICA) (2015) *Bases cartográficas contínuas, transporte*. http://geoftp.ibge.gov.br/cartas_e_mapas/bases_cartograficas_continuas/bc250/ [accessed 12 September 2018].
- KRAMER-SCHADT, S., NIEBALLA, J., PILGRIM, J.D., SCHRÖDER, B., LINDENBORN, J., REINFELDER, V. et al. (2013) The importance of correcting for sampling bias in *MaxEnt* species distribution models. *Diversity and Distributions*, 19, 1366–1379.
- KUKKALA, A.S. & MOILANEN, A. (2013) Core concepts of spatial prioritisation in systematic conservation planning. *Biological Reviews*, 88, 443–464.
- LEAL, I.R., SILVA, J.M.C., TABARELLI, M. & LACHER, JR, T.E. (2005) Changing the course of biodiversity conservation in the Caatinga of northeastern Brazil. *Conservation Biology*, 19, 701–706.
- MILLER, J. (2010) Species distribution modeling. *Geography Compass*, 4, 490–509.
- MMA (MINISTÉRIO DO MEIO AMBIENTE) (2016) *Sistema Nacional de Unidades Conservação—SNUC*. <http://www.mma.gov.br/areas-protegidas/sistema-nacional-de-ucs-snuc> [accessed 20 June 2017].
- MOILANEN, A., FRANCO, A.M.A., EARLY, R.I., FOX, R., WINTLE, B. & THOMAS, C.D. (2005) Prioritizing multiple-use landscapes for conservation: methods for large multi-species planning problems. *Proceedings of the Royal Society B*, 272, 1885–1891.
- MOILANEN, A., ANDERSON, B.J., EIGENBROD, F., HEINEMEYER, A., ROY, D.B., GILLINGS, S. et al. (2011) Balancing alternative land uses in conservation prioritization. *Ecological Applications*, 21, 1419–1426.
- MOILANEN, A., POUZOLS, F.M., MELLER, L., VEACH, V., ARPONEN, A., LEPPÄNEN, J. & KUJALA, H. (2017) *Spatial Conservation Planning Methods and Software: Zonation Version 4 User Manual*. Biodiversity Conservation Informatics Group, Department of Biosciences, University of Helsinki, Helsinki, Finland.
- OLIVEIRA, M.M. & KIERULFF, M.C.M. (2008) *Alouatta ululata*. In *The IUCN Red List of Threatened Species 2008: e.T918A13094890*. <http://dx.doi.org/10.2305/IUCN.UK.2008.RLTS.T918A13094890.en> [accessed 20 June 2017].
- OLIVEIRA, M.M., FERREIRA, J.G., MOTA, G.L.S. & SOARES, S.G. (2007) Mapeamento das áreas de ocorrência de *Alouatta ululata*—Etapa Ceará. In *A Primatologia do Brasil*, Volume 10 (ed. J.C. Bicca-Marques), pp. 151–161. Sociedade Brasileira de Primatologia, Porto Alegre, Brazil.
- PBMC (PAINEL BRASILEIRO DE MUDANÇAS CLIMÁTICAS) (2013) *Executive Summary: Scientific Basis of Climate Change—Contribution from the Working Group 1 to the First National Assessment Report of the Brazilian Panel on Climate Change (GT1 RAN1 PBMC)*. Universidade Federal do Rio de Janeiro, Rio de Janeiro, Brazil.
- PHILLIPS, S. (2008) *A Brief Tutorial on Maxent*. https://biodiversityinformatics.amnh.org/open_source/maxent/Maxent_tutorial2017.pdf [accessed 12 September 2018].
- PHILLIPS, S.J., ANDERSON, R.P. & SCHAPIRE, R.E. (2006) Maximum entropy modeling of species geographic distributions. *Ecological Modelling*, 190, 231–259.
- PRESSEY, R.L., CABEZA, M., WATTS, M.E., COWLING, R.M. & WILSON, K.A. (2007) Conservation planning in a changing world. *Trends in Ecology & Evolution*, 22, 583–592.
- PRESSEY, R.L., WATTS, M.E., BARRETT, T.W. & RIDGES, M.J. (2009). The C-Plan conservation planning system: origins, applications, and possible futures. In *Spatial Conservation Prioritization* (eds A. Moilanen, K.A. Wilson & H. Possingham), pp. 211–234. Oxford University Press, Oxford, UK.
- QGIS DEVELOPMENT TEAM (2016) *QGIS: A Free and Open Source Geographic Information System*. Open Source Geospatial Foundation. <http://www.qgis.org> [accessed 24 May 2017].
- RAINHO, A. & PALMEIRIM, J.M. (2013) Prioritizing conservation areas around multispecies bat colonies using spatial modeling. *Animal Conservation*, 16, 438–448.
- RILEY, S.J., DEGLORIA, S.D. & ELLIOT, R. (1999) A terrain ruggedness index that quantifies topographic heterogeneity. *Intermountain Journal of Sciences*, 5, 23–27.
- SDAT (SPATIAL DATA ACCESS TOOL) (2011) *Global 1km Forest Canopy Height*. https://webmap.ornl.gov/wcsdown/dataset.jsp?dg_id=10023_1 [accessed 12 September 2018].
- SEDAC (SOCIOECONOMIC DATA AND APPLICATIONS CENTER) (2011) *Global Rural-Urban Mapping Project (GRUMP), v1: Population Density Grid*. <http://dx.doi.org/10.7927/H4R2oZ93> [accessed 12 September 2018].
- SIMARD, M., PINTO, N., FISHER, J.B. & BACCINI, A. (2011) Mapping forest canopy height globally with spaceborne lidar. *Journal of Geophysical Research*, 116, 1–12.
- TORRES, R.R., LAPOLA, D.M. & GAMARRA, N.L.R. (2017) Future climate change in the Caatinga. In *Caatinga: The Largest Tropical Dry Forest Region in South America* (eds J.M.C. da Silva, I.R. Leal & M. Tabarelli), pp. 383–410. Springer, Cham, Switzerland.
- USGS (UNITED STATES GEOLOGICAL SURVEY) (2000) *Shuttle Radar Topography Mission 3 (SRTM3)*. <https://earthexplorer.usgs.gov> [accessed 12 September 2018].
- VIANA, M.C., BONVICINO, C.R., FERREIRA, J.G., JERUSALINSKY, L., LANGGUTH, A. & SEUÁNEZ, H. (2015) Understanding the relationship between *Alouatta ululata* and *Alouatta belzebul* (Primates: Atelidae) based on cytogenetics and molecular phylogeny. *Oecologia Australis*, 19, 173–182.
- WATTS, M.E., BALL, I.R., STEWART, R.S., KLEIN, C.J., WILSON, K., STEINBACK, C. et al. (2009) *Marxan with Zones*: software for optimal conservation based land- and sea-use zoning. *Environmental Modelling & Software*, 24, 1513–1521.
- WIENS, J.A., STRALBERG, D., JONGSOMJIT, D., HOWELL, C.A. & SNYDER, M.A. (2009) Niches, models, and climate change: assessing the assumptions and uncertainties. *Proceedings of the National Academy of Sciences of the United States of America*, 106, 19729–19736.
- ZOMER, R.J., TRABUCCO, A., BOSSIO, D.A. & VERCHOT, L.V. (2008) Climate change mitigation: a spatial analysis of global land suitability for clean development mechanism afforestation and reforestation. *Agriculture, Ecosystems & Environment*, 126, 67–80.